


ED Headache Treatment Algorithm

Orderset: ED headache



Reassessments should take place within one hour after each intervention.

Prochlorperazine (Compazine):

- Preferred first-line agent
- 0.15 mg/kg IV; max of 10 mg
- For patients with previous adverse reactions who refuse prochlorperazine, consider:
 - Metoclopramide (Reglan)
 - Give a half-dose of Prochlorperazine
 - Slow the infusion rate by half
 - Have patient walk during infusion if able and not dizzy to help with akathisia

Valproic Acid:

- Utilize “ED headache order” set to assure correct dosing and rate
- Patients who are discharged after treatment with valproic acid should be sent home on a 14-day course of oral valproic acid.
- < 10 years OR < 50 kg: Depakote ER 250 mg PO BID x 2 weeks
- >= 10 years AND >=50 kg: Depakote ER 500 mg PO BID x 2 weeks
- Give 1st oral dose in the ED
- If IV Valproic Acid unavailable do not give PO valproic acid in ED or send home with 14-day prescription

Dihydroergotamine (DHE) – See Lexicomp for additional details:

- Contraindications: (1) Triptan within 24 hours; (2) DHE within 14 days.
- MUST obtain pregnancy test in adolescent females
- < 30 kg: 0.5 mg; >= 30 kg: 1 mg
- Administer 50% of dose first (over 3 minutes) then remaining 50% in 30 minutes
- Expected Side Effects (counsel patient to anticipate): nausea, vomiting, flushing, and hypertension – most resolve within 1 hour so if patient has symptoms resolve and HA resolve, can go home
- After DHE in the ED, please discuss with neuro, as most patients will be admitted to neuro (or HM at LT after neuro discussion) unless HA at baseline and family prefers discharge, or family does not want further DHE
- If patient discharged home, advise no triptan for 24 hours after last dose of DHE

Steroids:

- Should be considered in children with recurrent headaches despite ED treatment (return visits within 72 hours for rebound headache)
 - Dexamethasone 0.6 mg/kg IV x1; max of 10 mg IV (No medications for discharge home)